


10/12/12

Test Notes

$T_p(M)$ is the set of all tangent vectors (not just unit tangents) to M at point p .

$\vec{c}(t)$ is a curve down on u .

#3



Assume that the directional derivatives are well-def.ncl.

#1-h We need to express \vec{x}_{11} in terms of \vec{x}_1, \vec{x}_2 , and \vec{n} .

Wrote

$$\vec{x}_{11} = -\vec{x}_1 + -\vec{x}_2 + -\vec{n}$$

and find the pieces!

$$\vec{c} : \mathbb{R} \rightarrow \mathbb{R}^2$$

$$\vec{x} : \mathbb{R}^2 \rightarrow \mathbb{R}^3$$

$$\vec{f} = \vec{x} \circ \vec{c} : \mathbb{R} \rightarrow \mathbb{R}^3$$

$$\text{Remember that } \vec{f}' = D\vec{x} \dot{\vec{c}}$$

Also remember that \vec{x}_i , \vec{x}_{ij} , \vec{n} are all also functions from U to \mathbb{R}^3 . We can therefore write something like $D\vec{n}$.

Let $\vec{f}(0) = p$ and $\vec{f} = \vec{x} \circ \vec{\varphi}$. There is some corresponding $q = \vec{c}(0) \in U$, such that $\vec{x}(q) = p$.

$$\vec{x}_i(q) = \vec{x}_i(c^1(0), c^2(0))$$


gives one particular tangent vector at p . \vec{x}_i is also a tangent vector for the u^1 curves.

#2 We are allowed to use k_n and k_g as we know them by definition.

Parallelism

"Intrinsic being" can measure the metric tensor. With (g_{ij}) we can measure length and angle.


Take the surface



$$2\pi R > \text{Circumference}$$



$$2\pi R \approx \text{C.r.c.}$$



This triangle has 270° !

We could also look at tangent vectors.

Let $V \subset M$. A vector field on V is a function

$$\vec{X} : V \rightarrow \mathbb{R}^3$$

tangent vector field : $\vec{X} : V \rightarrow T(M)$. So for
each p on V , $\vec{X}(p) \in T_p(M)$.

curve vector field : A tangent vector field along a curve $\vec{f} : [a, b] \rightarrow M$
is just a tangent vector field on $\vec{f} : [a, b]$.

Theorem :

A tangent vector field \vec{X} is parallel (to itself)
along \vec{f} iff $\frac{d\vec{X}}{dt}$ is parallel to \vec{n} .

This means that intrinsic beings couldn't measure the
~~length~~ change.

Equivalently, \vec{X}' always normal to the surface, or $\vec{X}' \perp \vec{n}$.

Proposition :

If $\vec{f} = \vec{x} \circ \vec{c}$ is a regular curve and $\vec{\Sigma}(t)$ is a v-f. on \vec{f} with $\vec{\Sigma}(t) = \sum \vec{x}^i(t) \vec{x}_i(c(t))$. We are simply expressing $\vec{\Sigma}$ in terms of \vec{x}_i and \vec{x}_j . the basis vectors for the tangent planes. Then $\vec{\Sigma}$ is parallel iff

$$\phi = \vec{\Sigma}^k + \sum \Gamma_{ij}^k \vec{x}^i \vec{c}^j$$

for $k=1, 2$.

Proof

$$\vec{\Sigma} = \sum \vec{x}^i \vec{x}_i$$

$$\dot{\vec{\Sigma}} = \sum \left(\vec{x}^i \vec{x}_i + \vec{x}^i \frac{d}{dt} \vec{x}_i \right)$$

Note that

$$\frac{d}{dt} \vec{x}_i = \frac{d}{dt} \vec{x}_i(u^1(t), u^2(t))$$

$$= \frac{\partial \vec{x}_i}{\partial u^1} \frac{du^1}{dt} + \frac{\partial \vec{x}_i}{\partial u^2} \frac{du^2}{dt}$$

$$= \cancel{\frac{\partial \vec{x}_i}{\partial u^1} \vec{c}^1} + \cancel{\frac{\partial \vec{x}_i}{\partial u^2} \vec{c}^2}$$

$$= \frac{\partial \vec{x}_i}{\partial u^1} \vec{c}^1 + \frac{\partial \vec{x}_i}{\partial u^2} \vec{c}^2$$

So

$$\dot{\vec{x}} = \sum_i \left(\dot{x}^i \vec{x}_i + \vec{x}^i \sum_j \dot{x}^j \Gamma_{ij}^k \vec{x}_k \right)$$

$\dot{\vec{x}}$ is parallel iff $\langle \dot{\vec{x}} | \vec{x}_j \rangle = 0$, for $j = 1, 2$.
 But

$$\langle \dot{\vec{x}} | \vec{x}_j \rangle = \sum_i \left(\dot{x}^i g_{jl} + \vec{x}^i \sum_j \dot{x}^j \Gamma_{ijl} \right) = 0$$

Sum against g^{lk} .

$$\sum_l \left(\sum_i \left(\dot{x}^i g_{jl} g^{lk} + \vec{x}^i \sum_j \dot{x}^j \Gamma_{ijl} g^{lk} \right) \right)$$

$$= \sum_i \left(\dot{x}^i \delta_i^k + \vec{x}^i \sum_j \dot{x}^j \Gamma_{ij}^k \right)$$

$$= \boxed{ \sum_i \left(\dot{x}^k + \sum_j \Gamma_{ij}^k \dot{x}^j \vec{x}^i \right) }$$

Theorem 6+7 :

Let \vec{y} be a curve (regular) on M and let $\tilde{\vec{x}} \in T_{\vec{y}(t_0)} M$.

Then there exists only one vector field \vec{x} such that
 \vec{x} is parallel along \vec{y} and $\vec{x}(t_0) = \tilde{\vec{x}}$.